



1 Types of Forces

Example 1.1

Weight on Moon = $mg = 20 \text{ kg} \times 1.6 \text{ m s}^{-2} = 32 \text{ N}$;

Weight on Mars = $mg = 20 \text{ kg} \times 3.7 \text{ m s}^{-2} = 74 \text{ N}$

Exercise 1.1

1.1 Gravitational Force

1. (a) Mass on Moon = $45 \text{ N} / 10 \text{ m s}^{-2} = 4.5 \text{ kg}$


(b) Weight on Moon = $4.5 \text{ kg} \times 1.6 \text{ m s}^{-2} = 7.2 \text{ N}$

2. Friction prevents them from moving towards each other.


3. $20 \text{ kg} \neq 200 \text{ N}$, because 20 kg is a mass while 200 N is weight (or a force). These are different types of physical quantities and hence cannot be equated with an equal sign. She should write the step as: weight = $20 \text{ kg} \times 10 \text{ m s}^{-2} = 200 \text{ N}$.

Example 1.2


(a)



(b)




(c)




block at rest on a ramp
(f prevents block from sliding down)

(d)




ball sliding to the right at constant speed

(e)



girl walking to the left
(f prevents foot from sliding to the right)



block accelerating to the right on a
conveyer belt (f prevents block from sliding to the left)

Exercise 2

2 Friction


4. (a) No. Horizontal force acts on ball by vertical wall (assume ball bounces off horizontally)
 (b) Yes. There is friction in the forward direction on each foot in contact with ground (see Example 3(b))
 (c) No. Drum skin vibrates perpendicular to itself (as it interacts with air)
 (d) No. Although there is significant friction (air resistance) opposing motion of rocket.

5. Draw arrow to represent friction **exactly** on the surface of contact.



(a) 1 N to left, not moving	(b) 2 N to left, not moving	(c) 3 N to right, not moving
(d) 5 N to left, not moving	(e) 5 N to left, accelerates to right	(f) No friction acting

6. Not necessary. There may be other forces that balanced the friction.

7.






Example 1.4




Exercise 1.4


1.4 Tension




10. tension acting downwards on the diving board




Example 1.5




A box resting on a surface



A man pushing against a wall




A ladder leaning against a wall




A cylinder resting on a slope

If the wall is rough, friction on the ladder acts upwards. The combined effect of friction and normal contact force acting on the ladder is represented by F_c .



2. Free-body diagram*


Example 2.1



Exercise 2

2 Free-body Diagram


1. (a) Normal contact force on ball by hand



(normal contact force balances weight)

(b)


Normal contact force on ball by hand



Weight of ball

(normal contact force > weight)

(c)













Weight of ball


(d) No. The force exerted by the hand is a contact force, which ceases to exist after the ball leaves the hand.

(e) No. It will move upwards but start to decelerate due to gravitation force acting on it.


2.

<p>(a) Static.</p> 	<p>(b) Static.</p> 
<p>(c) Static.</p> 	<p>(d) Static.</p> 
<p>(e) Rising vertically upwards</p> 	<p>(f) At the top of a vertical path.</p> 
<p>(g) Falling rock. Ignore air resistance.</p> 	<p>(h) Falling at constant speed.</p> 
<p>(i) Sliding to the right at constant speed without friction.</p> 	<p>(j) Moving to the right and decelerating due to friction.</p> 

3. (a) F_1 = normal contact force on A by B
 F_2 = normal contact force on B by A
 F_3 = normal contact force on B by surface



- (b) R_1 = normal contact force on A by surface



R_2 = normal contact force on B by surface
 T_1 = tension acting on A
 T_2 = tension acting on B
 W_A = weight of A
 W_B = weight of B
 f = friction

3 Balanced and Unbalanced forces

Example 3.1


The forces acting on it are unbalanced and the resultant force is 4 N to the left.

Exercise 3


3 Balanced and Unbalanced forces

1.


(a) Three horizontal forces being applied on a box of mass 2 kg at rest on a table.



(b) A submarine remains stationary below the sea.




(c) A plane of mass 50 000 kg moves through the air at constant speed.



4 Newton's First Law of Motion

Example 4.1

- (a) By Newton's first law, the passenger will continue to move forward with constant speed in a straight line when the bus comes to a stop.
- (b) When the roller-coaster suddenly turns right, the people will continue to move in a straight line (as shown below) according to Newton's first law. Hence they appear to swing to the left of their seats



Exercise 4

4 Newton's First Law of Motion

1. Assumptions made: the train is moving in a straight line, the train is closed and there is an absence of air resistance acting on the girl.
 - (a) train at constant velocity:

She will land on the same spot from which she jumped. Before jumping upward, she had a forward velocity, which is same as that of the train. When she jumped, she continue to move with the same forward velocity as the train, according to Newton's first law of motion.
 - (b) train with positive acceleration in a straight line:

She will land behind the spot from which she jumped. When she jumped, she continue to move with the same forward velocity but the train is speeding up.
 - (c) train with negative acceleration (slowing down) in a straight line:

She will land in front of the spot from which she jumped. When she jumped, she continue to move with the same forward velocity but the train is slowing down.
2. (a) A car is accelerating from rest: Accelerating means velocity is not constant. Hence, resultant force is not zero
- (b) The moon circling around the Earth: Change in direction of velocity means velocity if not constant, hence, there is resultant force.
- (c) An apple falling to the ground: Velocity is not constant means resultant force is not zero.
- (d) A car moving at constant speed around a corner: Change in direction of velocity means velocity if not constant, hence, there is resultant force.

5 Newton's Second Law of Motion

Example 5.1

(a) $F_{\text{net}} = ma = (0.120 \text{ kg})(0.60 \text{ m s}^{-2}) = 0.072 \text{ N}$


(b) $a = \frac{F_{\text{net}}}{m} = \frac{4.0 \text{ N}}{2.5 \text{ kg}} = 1.6 \text{ m s}^{-2}$

(c) $m = \frac{F_{\text{net}}}{a} = \frac{0.32 \text{ N}}{9.8 \text{ m s}^{-2}} = 0.03265 \text{ kg (4 s.f.)} \approx 0.033 \text{ kg (rounded off to 2 s.f.)}$

Example 5.2

(a) Resultant force $F_{\text{net}} = 50 \text{ N} - 10 \text{ N} = 40 \text{ N}$

(b) $F_{\text{net}} = ma$
 $a = \frac{F_{\text{net}}}{m} = \frac{40 \text{ N}}{5.0 \text{ kg}} = 8.0 \text{ m s}^{-2}$



Exercises 5

5 Newton's Second Law of Motion

1. (a) 120 N (b) 80 N (c) 243 N (or 240 N: 2 sf)
(d) 30 N, 50 N (e) 10 N, 50 N (f) 25 N, 50 N
2. (a) 150 N (b) 50 N (c) 100 N

Terminal Velocity

Example 5.2

(a) (i) Net force = $mg = 50 \text{ kg} \times 10 \text{ m s}^{-2} = 500 \text{ N}$ (air resistance is negligible)

Net acceleration = 10 m s^{-2}

(ii) Net force = $mg - R = 500 \text{ N} - 60 \text{ N} = 440 \text{ N}$

Net acceleration = net force \div m = $440 \text{ N} \div 50 \text{ kg} = 8.8 \text{ m s}^{-2}$

(iii) Net force = $mg - R = 500 \text{ N} - 475 \text{ N} = 25 \text{ N}$


Net acceleration = net force \div m = $25 \text{ N} \div 50 \text{ kg} = 0.5 \text{ m s}^{-2}$

(b) When acceleration is zero, net force = zero. Hence, air resistance = weight = 500 N


6. Newton's Third Law of Motion (Action & Reaction)

Example 6.1

(a)



(b)



Therefore, two pairs of action-reaction forces are related to a stone that is at rest on the ground.

Example 6.2

1. (a) $F_{\text{net}} = ma$

$$a = F \div m = 20 \text{ N} \div (2.0 \text{ kg} + 3.0 \text{ kg}) = \underline{4.0 \text{ m s}^{-2}}$$


(b) For the 3.0 kg mass, $T = m a = 3.0 \text{ kg} \times 4.0 \text{ m s}^{-2} = \underline{12 \text{ N}}$

OR for the 2.0 kg mass, net force = $20 - T = ma$; $T = 20 \text{ N} - 2.0 \text{ kg} \times 4.0 \text{ m s}^{-2} = \underline{12 \text{ N}}$


Exercise 6.6

6.6 Newton's Third Law of Motion


1.



2.




3.



4. Tension on both masses are equal (same string, frictionless pulley)
 16 kg mass will move down and 10 kg mass will move up.
 (same *acceleration* because they are connected by a string)

On 16 kg mass: $160 - T = 16a$
 On 10 kg mass: $T - 100 = 10a$
 Solving for T, $1600 - 10T = 16(T - 100)$
 $T = \underline{123 \text{ N}}$
 Therefore, $a = \underline{2.3 \text{ m s}^{-2}}$




7 Vector Addition

7.1 Unbalanced forces (Resultant force)

Example 7.1

- (a) Resultant force = $45 \text{ N} + 20 \text{ N} - 15 \text{ N}$
 $= 50 \text{ N}$ to the right.




- (b) Scale 1.0 cm represents 2.0 N

Magnitude of the resultant force = 17.2 N
 direction of resultant force: 13° anticlockwise from 11.0 N force

- (c) Scale 1.0 cm represents 100 N


Magnitude of the resultant force = 740 N
 Its direction is 24° west of north




Exercise 7.1

7.1 Unbalance forces (Resultant force)

1. Maximum resultant (sum of forces)



Minimum resultant (difference of forces)




2. Resultant force = 8700 N (± 100); Direction is 30° ($\pm 1^\circ$) anticlockwise from force Q.

7.2 Balanced forces (Object at equilibrium)

Example 7.2


Scale 1.0 cm represents 50 N

The magnitude of the third force F is 220 N and it acts 96° clockwise from the 150 N force.



Discussion :

Case on the right is incorrect. Extend R. Direction of T is 140° clockwise from R.



7.3 Comparison of vector triangles


Question: The resultant force R and the 3rd force F_3 are in opposite directions.

Exercise 7.2


7.2 Balanced forces (Object at equilibrium)

1. (a) (i) Scale: 1.0 cm represents 5.0 N
 T (5.0 cm) = 25 N
 (ii) Since vector triangle is an equilateral triangle,
 $T = \underline{25 N}$.


- (b) The tensions listed in order from highest to the lowest is T_3, T_1, T_2



2. (a)



- (b)



- (c) T_1 is greater than T_2 as T_1 although T_1 has a shorter string than T_2 . Therefore, a longer string does not imply a greater amount of tension.
- (d) Since the angles are equal, the vector diagram for blocks A and B are the same! The tensions, $T_1=T_3$ and $T_2=T_4$. The tensions are dependent on the angles of inclination of the strings and not the length of the strings.
- (e) No. The triangle has a horizontal side but the weight is vertical!

8 Vector Resolution of Forces

Example 8.1


(a) $F_x = 320 \cos 40^\circ = \underline{245 \text{ N}}$

(b) $F_y = 320 \sin 40^\circ = \underline{206 \text{ N}}$

Example 8.2

(a) $W_x = W \sin \theta$

(a) $W_y = W \cos \theta$



Question: The original triangle has a horizontal side, but the weight is vertical!

Example 8.3

$QR_x = PQ = QR \cos 30^\circ$

$QR_y = 10 \text{ N} = QR \sin 30^\circ$

---> $PQ/10 = \cos 30^\circ / \sin 30^\circ$

$PQ = 10 / \tan 30^\circ$

$= 17.3$

$= 17 \text{ N (2 sf)}$

Exercise 8

8 Vector Resolution of Forces

1. (a) Horizontal component = $25 \cos 40^\circ = \underline{19 \text{ N}}$

(b) Net force = horizontal component = $19 \text{ N} = ma$; $\therefore a = 19 \text{ N} \div 2.0 \text{ kg} = \underline{9.5 \text{ m s}^{-2}}$