



2024 Sec 4 Advanced Physics Exercise 2A

Electric Fields

Name: _____ () Class: 4/ ___ Date: _____

Given formulae and constant:

$$E = \frac{F}{q}$$

$$F = \left| \frac{kQq}{r^2} \right|$$

where $k = 8.99 \times 10^9 \text{ kg m}^3 \text{ s}^{-4} \text{ A}^{-2}$

$$v = r \omega$$

$$a = \frac{v^2}{r}$$

For relevant calculations, you may take the mass of a proton to be $1.67 \times 10^{-27} \text{ kg}$ and its charge to be $1.60 \times 10^{-19} \text{ C}$.

A Two identical point charges were initially separated by a distance of d . Due to the mutual repulsion, they move away from each other. When the separation increased by 5.0 mm, the repulsion force decreased from $100 \mu\text{N}$ to $25 \mu\text{N}$.

(a) Calculate the value of d .

$d = \dots\dots\dots$


(b) Determine the magnitude of the charges.

magnitude of charges = $\dots\dots\dots$

- B** A proton travelling along the x-axis is slowed down by a uniform electric field E . At $x = 30$ cm, the proton has a speed of 2.5×10^6 m s⁻¹ and at $x = 60$ cm, its speed is zero. Find the magnitude and direction of E .

$$E = \dots\dots\dots$$

- C** A beam of protons are projected with an initial speed of $v_i = 9.55 \times 10^3$ m s⁻¹ into a region of uniform electric field $E = 720$ N C⁻¹ directed vertically downwards. The protons are to hit a target that lies at a horizontal distance of 1.27 mm away from the point where the protons cross the plane and enter the electric field.



- (a)** Determine the two possible angles θ_1 and θ_2 that will result in a successful hit. Ignore gravitational acceleration.

$$\theta_1 = \dots\dots\dots$$

$$\theta_2 = \dots\dots\dots$$

- (b) Determine the time interval during which the proton is above the plane and within the region of the electric field for each trajectory.

$t_1 = \dots\dots\dots$

$t_2 = \dots\dots\dots$

$E = \dots\dots\dots$

Answers:	1(a). 5.0 mm	1(b). 5.3×10^{-10} C	2. 138.1 g	3. -1.09×10^5 N C ⁻¹
	4(a). 36.9° / 53.1°	4(b). 166 ns / 221 ns	5(a). out of the page	
	5(b). 0.39 m	5(c)(i). upwards	5(c)(ii). 5.4×10^5 N C ⁻¹	